

Benders Decomposition

Operations Research

Anthony Papavasiliou

- 1 Cutting Plane Methods
- 2 Context and Description of Benders Decomposition
- 3 Useful Results
- 4 Statement of Algorithm and Proof of Convergence
- 5 Example: Capacity Expansion Planning

Table of Contents

- 1 Cutting Plane Methods
- 2 Context and Description of Benders Decomposition
- 3 Useful Results
- 4 Statement of Algorithm and Proof of Convergence
- 5 Example: Capacity Expansion Planning

Cutting plane methods: optimization methods which are based on the idea of iteratively refining the objective function or set of feasible constraints of a problem through linear inequalities

Kelley's Cutting Plane Algorithm

Kelley's cutting plane algorithm is designed for solving convex non-differentiable optimization problems:

$$\begin{aligned} z^* &= \min c^T x + F(x) \\ \text{s.t. } x &\in X \end{aligned}$$

where

- X is a compact convex subset of \mathbb{R}^n
- $F : \mathbb{R}^n \rightarrow \mathbb{R}$ is a convex function
- $c \in \mathbb{R}^n$ is a parameter vector

Kelley's Cutting Plane Algorithm

Define

- $L_k : \mathbb{R}^n \rightarrow \mathbb{R}$ as lower bounding function of $F(x)$ at iteration k
- Lower bound L_k of z^* at iteration k
- Upper bound U_k of z^* at iteration k

Idea: gradually bound $F(x)$ from below with functions $L_k(x)$

Kelley's Cutting Plane Algorithm

Step 0: Set $k = 0$, and assume $x_1 \in X$ given. Set $L_0(x) = -\infty$ for all $x \in X$, $U_0 = c^T x_1 + F(x_1)$, and $L_0 = -\infty$

Step 1: Set $k = k + 1$. Find $a_k \in \mathbb{R}$ and $b_k \in \mathbb{R}^n$ such that

$$F(x_k) = a_k + b_k^T x_k$$

$$F(x_k) \geq a_k + b_k^T x, x \in X$$

Step 2: Set

$$U_k = \min(U_{k-1}, c^T x_k + F(x_k))$$

and

$$L_k(x) = \max(L_{k-1}(x), a_k + b_k^T x), x \in X$$

Step 3: Compute

$$L_k = \min_{x \in X} c^T x + L_k(x)$$

and denote x_k as the optimal solution of this problem

Step 4: If $U_k - L_k = 0$, stop; else, repeat from step 1

Nomenclature of Cutting Plane Methods

- **Benders decomposition**: specific method for obtaining the cutting planes when $F(x)$ is the value function of a second-stage linear program
- **L-shaped method**: specific instance of Benders decomposition when second-stage linear program is decomposable into a set of scenarios
- **Multi-cut L-shaped method**: alternative to L-shaped method which generates multiple cutting planes at step 1 of Kelley's method
- Cutting plane methods generalized to **bundle methods** in non-differentiable convex optimization (commonly used in Lagrange relaxation)

Table of Contents

- 1 Cutting Plane Methods
- 2 Context and Description of Benders Decomposition**
- 3 Useful Results
- 4 Statement of Algorithm and Proof of Convergence
- 5 Example: Capacity Expansion Planning

When to Use Benders Decomposition

Consider the following optimization problem:

$$z^* = \min c^T x + q^T y$$

$$Ax = b$$

$$Tx + Wy = h$$

$$x, y \geq 0$$

with $x \in \mathbb{R}^{n_1}$, $y \in \mathbb{R}^{n_2}$, $c \in \mathbb{R}^{n_1}$, $b \in \mathbb{R}^{m_1}$, $A \in \mathbb{R}^{m_1 \times n_1}$, $q \in \mathbb{R}^{n_2}$,
 $h \in \mathbb{R}^{m_2}$, $T \in \mathbb{R}^{m_2 \times n_1}$, $W \in \mathbb{R}^{m_2 \times n_2}$

- This is not (necessarily) a stochastic program
- This is a two-stage program

Context for Benders decomposition:

- 1 entire problem is difficult to solve
- 2 if $Tx + Wy = h$ is ignored, problem is relatively easy
- 3 if x is fixed, problem is relatively easy

Idea of Benders Decomposition

Define **value function** $V : \mathbb{R}^{n_1} \rightarrow \mathbb{R}$

$$(S) : \quad V(x) = \min_y q^T y \\ Wy = h - Tx \\ y \geq 0$$

Equivalent description of problem

$$\min c^T x + V(x) \\ Ax = b \\ x \in \text{dom } V \\ x \geq 0$$

Note: $\text{dom } V = \{x : \exists y, Tx + Wy = h, y \geq 0\}$

Graphical Description of Benders Decomposition




Table of Contents

- 1 Cutting Plane Methods
- 2 Context and Description of Benders Decomposition
- 3 Useful Results**
- 4 Statement of Algorithm and Proof of Convergence
- 5 Example: Capacity Expansion Planning


Dual of Second-Stage Linear Program

The dual of (S) can be expressed as:

$$(D) : \max_{\pi} \pi^T (h - Tx)$$
$$\pi^T W \leq q^T$$

Note: feasible region of (D) does *not* depend on x

- V : set of extreme points of $\pi^T W \leq q^T$
- R : set of extreme rays of $\pi^T W \leq q^T$



$\pi \in V, \sigma \in R$ do *not* depend on x , can be enumerated

Value Function Is Piecewise Linear

- $V(x)$ is a piecewise linear convex function of x
- If π_0 is dual optimal multiplier of (S) given x_0 , then

$$\pi_0^T(h - Tx_0)$$

is a supporting hyperplane of $V(x)$ at x_0

We recall a previous result for the proof

Parametrizing the Right-Hand Side

Define $c(u)$ as optimal value of

$$c(u) = \min f_0(x)$$
$$f_i(x) \leq u_i, i = 1, \dots, m$$


where $x \in \text{dom } f_0$ is the convex domain of $f_0(x)$ and f_0, f_i are convex functions

- $c(u)$ is convex
- Suppose strong duality holds and denote λ^* as the maximizer of the dual function $\inf_{x \in \text{dom } f_0} (f_0(x) - \lambda^T (f(x) - u))$ for $\lambda \leq 0$. Then $\lambda^* \in \partial c(u)$.



From previous result:

- $V(h - Tx)$ is convex, so $V(x)$ is convex
- $\pi_0 \in \partial V(h - Tx_0)$, so $\pi_0^T(h - Tx)$ is a supporting hyperplane of $V(x)$ at x_0
- (S) has a finite number of dual optimal multipliers \Rightarrow finite number of supporting hyperplanes for $V(x) \Rightarrow V(x)$ is piecewise linear convex



Domain of Value Function

dom V can be expressed equivalently as follows:

$$\text{dom } V = \{\sigma^T(h - Tx) \leq 0, \sigma \in R\}$$


where $\sigma \in R$ is the set of extreme rays of $\pi^T W \leq q^T$

Proof that $\text{dom } V \subseteq \{\sigma^T(h - Tx) \leq 0, \sigma \in R\}$:

- Suppose $x \in \text{dom } V$ and $\sigma^T(h - Tx) > 0$ for some $\sigma \in R$
- σ is an extreme ray $\Rightarrow \sigma^T W \leq 0$
- Consider any dual feasible vector π_0 : $\pi_0 + \lambda\sigma$ is feasible for any $\lambda \geq 0$
- Since $\sigma^T(h - Tx) > 0$, (D) becomes unbounded
- Contradiction with assumption that $x \in \text{dom } V \Rightarrow \sigma^T(h - Tx) \leq 0$ for all $\sigma \in R$

Proof that $\{\sigma^T(h - Tx) \leq 0, \sigma \in R\} \subseteq \text{dom } V$:

- Any ray of $\pi^T W \leq q^T$ can be expressed as convex combination of extreme rays
- Therefore, for any ray σ of $\pi^T W \leq q^T$ it follows that $\sigma^T(h - Tx) \leq 0 \Rightarrow (D)$ cannot become unbounded



$$\min c^T x + \theta$$

$$Ax = b$$

$$\sigma_r^T (h - Tx) \leq 0, \sigma_r \in R$$

$$\theta \geq \pi_v^T (h - Tx), \pi_v \in V$$

$$x \geq 0$$

θ : free auxiliary variable

Relax inequalities that define $V(x)$ and $\text{dom } V$:

$$(M) : z_k = \min c^T x + \theta$$


$$Ax = b$$

$$\sigma^T (h - Tx) \leq 0, \sigma \in R_k \subseteq R$$

$$\theta \geq \pi^T (h - Tx), \pi \in V_k \subseteq V$$

$$x \geq 0$$

Bounds and Exchange of Information



Solution of master problem provides:

- lower bound $z_k \leq z^*$
- candidate solution x_k
- under-estimator of $V(x_k)$, $\theta_k \leq V(x_k)$

Solution of slave problem with input x_k provides:

- upper bound $c^T x_k + q^T y_{k+1} \geq z^*$
- new vertex π_{k+1} or new extreme ray σ_{k+1}

Table of Contents

- 1 Cutting Plane Methods
- 2 Context and Description of Benders Decomposition
- 3 Useful Results
- 4 Statement of Algorithm and Proof of Convergence**
- 5 Example: Capacity Expansion Planning

Benders Decomposition Algorithm

Step 0: Set $k = 0$, $V_0 = R_0 = \emptyset$.

Step 1: Solve (M) . Store x_k .

- If (M) is feasible, store x_k .
- If (M) is infeasible, exit. Problem is infeasible.

Step 2: Solve (S) with x_k as input.

- If (S) is infeasible, let $R_{k+1} = R_k \cup \{\sigma_{k+1}\}$. Let $k = k + 1$ and return to step 1.
- If (S) is feasible, let $V_{k+1} = V_k \cup \{\pi_{k+1}\}$
 - If $V_k = V_{k+1}$, terminate with (x_k, y_{k+1}) as optimal solution.
 - Else, let $k = k + 1$ and return to step 1.

Finite termination since V and R are finite

Proof of Convergence

Denote x_k as solution of (M) and use it as input in (S)

- Suppose (S) is feasible, denote π_{k+1} as optimal vertex. If $\pi_{k+1} \in V_k$ then x_k is optimal.
- Suppose (S) is infeasible, denote σ_{k+1} as extreme ray. Then $\sigma_{k+1} \notin R_k$.

Proof that $\pi_{k+1} \in V_k \Rightarrow x_k$ is optimal

- For any x feasible, $c^T x + V(x) \geq c^T x_k + \theta_k$ because (M) is a relaxation of the original problem
- If $\theta_k = V(x_k)$, then x_k is optimal since for any x feasible, $c^T x + V(x) \geq c^T x_k + V(x_k)$
- We already know that $\theta_k \leq V(x_k)$ (first bullet)
- Need to show that $\theta_k \geq V(x_k)$ (next slide)

Proof that $\pi_{k+1} \in V_k \Rightarrow \theta_k \geq V(x_k)$

- We know that $V(x_k) = \pi_{k+1}^T (h - Tx_k)$ (why?)
- Since $\theta \geq \pi^T (h - Tx)$, $\pi \in V_k$ is enforced in (M) at iteration k , if $V_{k+1} = V_k$ then $\theta_k \geq \pi_{k+1}^T (h - Tx_k)$
- Combining the above relationships,
$$\theta_k \geq \pi_{k+1}^T (h - Tx_k) = V(x_k)$$


Proof that (S) infeasible $\Rightarrow \sigma_{k+1} \notin R_k$

- σ_{k+1} is an extreme ray $\Rightarrow \sigma_{k+1}^T (h - Tx_k) > 0$
- If $\sigma_{k+1} \in R_k$, then $\sigma_{k+1}^T (h - Tx_k) \leq 0$ (contradicting the first bullet)
- Therefore, $\sigma_{k+1} \notin R_k$

Table of Contents

- 1 Cutting Plane Methods
- 2 Context and Description of Benders Decomposition
- 3 Useful Results
- 4 Statement of Algorithm and Proof of Convergence
- 5 Example: Capacity Expansion Planning**

Load Duration Curve



Load duration curve is obtained by sorting load time series in descending order

Mathematical Programming Formulation

$$\min_{x, y \geq 0} \sum_{i=1}^n (l_i \cdot x_i + \sum_{j=1}^m C_i \cdot T_j \cdot y_{ij})$$

$$\text{s.t. } \sum_{i=1}^n y_{ij} = D_j, j = 1, \dots, m$$

$$\sum_{j=1}^m y_{ij} \leq x_i, i = 1, \dots, n-1$$

- l_i, C_i : fixed/variable cost of technology i
- D_j, T_j : height/width of load block j
- y_{ij} : capacity of i allocated to j
- x_i : capacity of i

Problem Data

Technology	Fuel cost (\$/MWh)	Inv cost (\$/MWh)
Coal	25	16
Gas	80	5
Nuclear	6.5	32
Oil	160	2

	Duration (hours)	Level (MW)
Base load	8760	0-4235
Medium load	7000	4235-7496
Peak load	1500	7496-10401

$$(M) : \min_{x \geq 0} \sum_{i=1}^n l_i \cdot x_i + \theta$$
$$\theta \geq \sum_{j=1}^m \lambda_j^v D_j + \sum_{i=1}^n \rho_i^v x_i, (\lambda^k, \rho^k) \in V_k$$
$$\theta \geq 0$$

λ_j^k, ρ_i^k : dual optimal multipliers of slave

Note $\theta \geq 0$

- because slave has non-negative cost
- necessary for boundedness of master

$$(S) : \min_{y \geq 0} \sum_{i=1}^n \sum_{j=1}^m C_i \cdot T_j \cdot y_{ij}$$

$$(\lambda_j) : \sum_{i=1}^n y_{ij} = D_j, j = 1, \dots, m$$

$$(\rho_i) : \sum_{j=1}^m y_{ij} \leq \bar{x}_i, i = 1, \dots, n-1$$

\bar{x}_i : trial decision from master

Sequence of Investments

Iteration	Coal (MW)	Gas (MW)	Nuclear (MW)	Oil (MW)
1	0	0	0	0
2	0	0	0	8735.6
3	0	0	0	18565.1
4	0	14675.8	0	0
5	10673.3	0	0	0
6	0	0	7337.9	3063.1
7	0	1497.7	7337.9	732.2
8	0	1497.7	7337.9	2033.3
9	0	0	8966	1435
10	2851.8	2187.2	5362	0
11	8321	0	0	2080
12	6989.5	4489.5	56.5	0
13	3261	2905	4235	0

- A *new* investment proposal is necessarily made in each iteration (why?)
- Greedy behavior
 - First iteration: no investment
 - Early iterations: technologies with low investment cost