

Reviews of Probability Theory and Convex Analysis

Operations Research

Anthony Papavasiliou

1 Probability

2 Convex Analysis

Table of Contents

1 Probability

2 Convex Analysis


Given a **sample space** Ω , a sigma algebra **sigma-algebra** \mathcal{A} is a set of subsets of Ω such that

- $\Omega \in \mathcal{A}$
- if $A \in \mathcal{A}$ then also $\Omega - A \in \mathcal{A}$
- if $A_i \in \mathcal{A}$ for $i = 1, 2, \dots$ then also $\bigcup_{i=1}^{\infty} A_i \in \mathcal{A}$

Sigma Algebras for Markov Decision Processes

- Given a sample space Ω , there is no unique sigma-algebra of Ω , here are two
 - $\{\emptyset, \Omega\}$
 - 2^Ω set of all subsets (**power set**) of Ω
- In these notes we will focus on finite Ω , and its power set, denoted $\mathcal{B}(\Omega)$
- Elements of a sigma-algebra are called **events**

Example: Stock Price Evolution



- State space is the set of values that the stock price can take at each stage: $S_0 = \{C\}$, $S_1 = \{C_u, C_d\}$,
 $S_2 = \{C_{uu}, C_{ud}, C_{dd}\}$
- Sample space is

$$\Omega = S_0 \times S_1 \times S_2 = \{(C, C_u, C_{uu}), (C, C_u, C_{ud}), (C, C_u, C_{dd}), (C, C_d, C_{uu}), (C, C_d, C_{ud}), (C, C_d, C_{dd})\}$$

Information in period 2:

$$\begin{aligned}\Omega &= \{(C, C_u, C_{uu}), (C, C_u, C_{ud}), (C, C_d, C_{ud}), (C, C_d, C_{dd})\} \\ \mathcal{B}(\Omega) &= \{\emptyset, \{(C, C_u, C_{uu})\}, \dots, \{(C, C_u, C_{uu}), (C, C_u, C_{ud})\}, \dots, \\ &\{(C, C_u, C_{uu}), (C, C_u, C_{ud}), (C, C_d, C_{ud})\}, \dots, \\ &\{(C, C_u, C_{uu}), (C, C_u, C_{ud}), (C, C_u, C_{dd}), (C, C_d, C_{uu})\}, \dots, \\ &\dots \\ &\{(C, C_u, C_{uu}), (C, C_u, C_{ud}), (C, C_u, C_{dd}), \\ &(C, C_d, C_{uu}), (C, C_d, C_{ud}), (C, C_d, C_{dd})\}\end{aligned}$$

'the stock price in period 2 is C_{ud} ': identifiable (corresponds to $\{(C, C_u, C_{ud}), (C, C_d, C_{ud})\}$, which is an element of $\mathcal{B}(\Omega)$)

Information in period 0:

$$\mathcal{A}_0 = \{\emptyset, \Omega\}.$$

This is a valid sigma-algebra on Ω (satisfies all three conditions of the definition of a sigma-algebra)

Information in period 1:

$$\begin{aligned} \mathcal{A}_1 = & \{\emptyset, \\ & \{(C, C_u, C_{uu}), (C, C_u, C_{ud}), (C, C_u, C_{dd})\}, \\ & \{(C, C_d, C_{uu}), (C, C_d, C_{ud}), (C, C_d, C_{dd})\}, \\ & \Omega\} \end{aligned}$$

- ‘the stock price in period 0 is C , and in period 1 it is C_u ’: distinguishable (2nd element in \mathcal{A}_1)
- ‘the stock price in period 0 was C , in period 1 it is C_u , and in period 2 it is C_{uu} ’: not distinguishable (not in \mathcal{A}_2)

A **measurable probability space** is the triplet $(\Omega, \mathcal{A}, \mathbb{P})$, where Ω is the sample space, \mathcal{A} is a sigma-algebra of Ω , and $\mathbb{P} : \mathcal{A} \rightarrow [0, 1]$ is the probability measure that obeys the following properties:

- $\mathbb{P}(\emptyset) = 0$,
- $\mathbb{P}(\Omega) = 1$, and
- $\mathbb{P}(\cup_{i=1}^{\infty} A_i) = \sum_i P(A_i)$ if A_i are disjoint

Note: $\mathbb{P}[\cdot]$ and $\mathbb{P}(\cdot)$ will be used interchangeably

A **random variable** $\xi : \Omega \rightarrow \mathbb{R}$ is a function that maps random outcomes to real values

A **random vector** is a function $\xi : \Omega \rightarrow \mathbb{R}^n$ that maps outcomes to real-valued vectors

Given an index set T , and a probability space $(\Omega, \mathcal{B}(\mathcal{A}), \mathbb{P})$, a **stochastic process** is a collection of \mathbb{R}^n -valued random vectors, which can be written as $(X(t) : t \in T)$

Given $(\Omega, \mathcal{B}(\Omega))$, a **filtration** is an increasing sequence of sigma-algebras $\{\mathcal{A}_t\}_{t \geq 0}$ where each t is non-negative and

$$t_1 \leq t_2 \Rightarrow \mathcal{A}_{t_1} \subseteq \mathcal{A}_{t_2}$$

In the stock price example, the sequence $(\mathcal{A}_0, \mathcal{A}_1, \mathcal{A}_2)$, where $\mathcal{A}_2 = \mathcal{B}(\Omega)$, defines a filtration on $(\Omega, \mathcal{B}(\Omega))$

The **conditional probability** of event A given event B is defined as

$$\mathbb{P}[A|B] = \begin{cases} \frac{\mathbb{P}[A \cap B]}{\mathbb{P}[B]}, & \mathbb{P}[B] > 0 \\ 0, & \mathbb{P}[B] = 0 \end{cases}$$

Stock Pricing Example

Random variables in stock pricing example: price ξ_t in stage t

For period 0,

$$\xi_0(\omega) = C, \omega \in \Omega$$

For period 1,

$$\xi_1(\omega) = C_U, \omega = (C, C_U, \cdot)$$

$$\xi_1(\omega) = C_D, \omega = (C, C_D, \cdot)$$

For period 2,

$$\xi_2(\omega) = C_{UU}, \omega = (C, \cdot, C_{UU})$$

$$\xi_2(\omega) = C_{UD}, \omega = (C, \cdot, C_{UD})$$

$$\xi_2(\omega) = C_{UD}, \omega = (C, \cdot, C_{UD})$$

Distribution Functions

The **cumulative distribution function** of a random variable ξ is defined as $F_\xi(x) = P(\xi \leq x)$

For discrete random variables, the **probability mass function** f is defined as $f(\xi^k) = P(\xi = \xi^k)$, $k \in K$ with $\sum_{k \in K} f(\xi^k) = 1$

For continuous random variables, the **density function** f is defined by $P(a \leq \xi \leq b) = \int_a^b f(\xi) d\xi = \int_a^b dF(\xi)$ with $\int_{-\infty}^{\infty} dF(\xi) = 1$

The **expectation** of a random variable is defined as

$\mu = \sum_{k \in K} \xi^k f(\xi^k)$ for discrete random variables and as $\int_{-\infty}^{\infty} \xi dF(\xi)$ continuous random variables

The moment **rth moment** of ξ is $\bar{\xi}^{(r)} = \mathbb{E}[\xi^r]$

The **variance** of a random variable is defined as $\mathbb{E}[(\xi - \mu)^2]$

The **α -quantile** of ξ is a point η such that for $0 < \alpha < 1$,

$$\eta = \min\{x | F(x) \geq \alpha\}$$

Convergence in Distribution

A sequence X_1, X_2, \dots of random variables is said to **converge in distribution**, or **converge weakly**, or **converge in law** to a random variable X if

$$\lim_{n \rightarrow \infty} F_n(x) = F(x)$$

for every $x \in \mathbb{R}$ at which F is continuous. F and F_n are the cumulative distribution functions of X and X_n respectively

Table of Contents

1 Probability

2 Convex Analysis



A **polyhedron** is a set in \mathbb{R}^n which can be expressed as $\{x : Ax \leq b\}$, where $A \in \mathbb{R}^m \times \mathbb{R}^n$ and $b \in \mathbb{R}^m$.

Convex

Consider a set of points $x_i \in \mathbb{R}^n, i = 1, \dots, n$, a **convex combination** of these points is a point $\sum_{i=1}^n \lambda_i x_i$, such that $\sum_{i=1}^n \lambda_i = 1$ and $\lambda_i \geq 0, i = 1, \dots, n$

X is a **convex set** if it contains any convex combination of points $x_i \in X$

The **convex hull** of a set of points is the set of all convex combinations of these points



Extreme Points, Extreme Rays

An **extreme point** of a convex set is a point which cannot be expressed as the convex combination of two distinct points in the set


A point $r \in \mathbb{R}^n$ is a **ray** of a polyhedron P if and only if for any point $x \in P$, $\{y \in \mathbb{R}^n : y = x + \lambda r, \lambda \geq 0\} \subseteq P$

A ray r of P is an **extreme ray** if it cannot be expressed as a convex combination of other rays of P

Convex and Concave Functions

f is a **convex function** if for all $0 \leq \lambda \leq 1$ and any x_1, x_2 we have $f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2)$

f is **concave** if $-f$ is convex.



Frequently Encountered Classes of Functions

f is an **additively separable function** if it can be written as

$$f(x_1, x_2, \dots, x_n) = f_1(x_1) + f_2(x_2) + \dots + f_n(x_n)$$

The **domain** of f , $\text{dom } f$, is the set where f is finite

A continuous function f is **piecewise linear** if it can be written as

$$f(x) = \max_{i=1, \dots, n} (a_i^T x + b_i)$$

for all $x \in \text{dom } f$, where $a_i \in \mathbb{R}^n$, $b_i \in \mathbb{R}$, and n a finite integer number

Convex Optimization Problems

An **optimization problem** is the problem of finding the minimum of a function f over a set $X \subset \mathbb{R}^n$:

$$\begin{aligned} \min f(x) \\ \text{subject to } x \in X \end{aligned}$$

X is the **feasible set** of the problem, f is the **objective function** of the problem

Any $x \in X$ is a **feasible solution**, any $x^* \in X$ such that $f(x^*) \leq f(x)$ for any $x \in X$ is an **optimal solution**

A **convex optimization problem** is an optimization problem with a convex objective function and a convex set of feasible solutions